



ferti-lome®

Caterpillar Killer with Bt

BIOLOGICAL INSECTICIDE



Controls Worms And Caterpillars
On Fruits, Vegetables,
Ornamentals And Shade Trees



ACTIVE INGREDIENT:

Bacillus thuringiensis subspecies *kustaki* strain SA-12 solids, spores and Lepidopteran active toxins (At least 6 million viable spores per mg of Active Ingredient)* 98.35%

OTHER INGREDIENTS: 1.65%

TOTAL: 100.00%

*The percent active ingredient does not indicate product performance and potency measurements are not federally standardized.

**KEEP OUT OF REACH
OF CHILDREN**
CAUTION

NET CONTENTS 8 FL. OZS. / 236 ML

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Harmful if inhaled. Avoid breathing vapors or spray mist. Remove and wash contaminated clothing before reuse.

FIRST AID

If in eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled:	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Hot Line Number: 1-800-255-3924.	

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. This product must not be applied within 300 feet of any habitats of threatened or endangered Lepidoptera (i.e., moths or butterflies) or Coleoptera (i.e., beetles).

Manufactured For:



230 FM 87 BONHAM, TEXAS 75418
EPA Reg. No. 70051-106-7401
EPA Est. No. 7401-TX-01



7 32221 16011 8

PRODUCT INFORMATION

ferti•lome® Caterpillar Killer with Bt is an easy to mix liquid concentrate. Properties include:

- Kills worms and caterpillar stage insects, but has no effect on birds, earthworms, or beneficial insects, such as honeybees and ladybugs, when used as directed.
- Worms and caterpillars eat treated foliage then immediately stop feeding and damaging plants.
- Acceptable for use on edible plants up to the day of harvest.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Not for use on agricultural establishments covered by the WPS (40 CFR Part 170).

Apply **ferti•lome® Caterpillar Killer with Bt** when worms or caterpillars are first noticed, then repeat at five (5) to seven (7) day intervals while they are active. Apply more frequently to control heavy infestations. Apply thoroughly to top and bottom of foliage. Reapply after heavy rains.

ferti•lome® Caterpillar Killer with Bt must be eaten by worms or caterpillars to be effective. After ingesting the insecticide, they immediately stop feeding, though they may otherwise appear to be unaffected for several days. Make applications when worms are small. They must be actively feeding treated, exposed foliage. You may apply **ferti•lome® Caterpillar Killer with Bt** up to and on the day of harvest.

HOW TO APPLY

Always shake or stir **ferti•lome® Caterpillar Killer with Bt** thoroughly before use. Partially fill sprayer with water before adding the application rate amount of product thoroughly and add remaining amount of water.

Spray leaf surfaces thoroughly, top and bottom, for complete control. Agitate regularly while spraying. Use all of spray mixture within 24 hours.

FRUITS AND VEGETABLES

Mix 1.5 to 2.0 fluid ounces per 3 gallons of water (3 to 4 teaspoons per gallon) for a hand sprayer and apply to all plant foliage to cover an area of 1000 square feet. Apply at first sign of infestation and repeat at weekly intervals when needed to maintain control of the following insects:

Cabbage looper on broccoli, cauliflower, collards, kale, mustard greens, turnip greens, cabbage, celery, lettuce, melons, and tomatoes.

Imported cabbage worm on broccoli, cabbage, cauliflower, collards, kale, mustard greens, and turnip greens.

Tomato Hornworm on tomatoes.

SHADE TREES AND ORNAMENTALS

Mix 2 fluid ounces per 3 gallons of water (4 teaspoons per gallon) for a hand sprayer or proportional amounts for a hose-end sprayer following manufacturer's directions. Apply at first sign of infestation and repeat at weekly intervals when needed to maintain control.

Apply to thoroughly cover all foliage surfaces of shade trees and ornamentals for control of the following leaf-feeding worms:

Bagworm	Spring cankerworm
Fall cankerworm	Gypsy moth*
Tent caterpillar	Elm spanworm
Fall webworm	

*Apply when buds swell, but before leaves are fully open. If eggs hatch over a long period of time or if reinfestation occurs, spray about 14 days after first application.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Pesticide Storage: Store in a cool dry place. Activity may be impaired by storage at temperatures over 90° F.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

Nonrefillable container. Do not reuse or refill this container. Never place unused product down any indoor or outdoor drain. For containers less than or equal to 5 gallons:

If Empty: Place in trash or offer for recycling if available.

If Partly Filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions.

Voluntary Purchasing Groups, Inc. warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the insect problem, condition of the crop, incompatibility with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OF MERCHANTABILITY IS MADE.

- Kills Bagworms, Fall Cankerworms, Tent Caterpillar, Fall Webworm, Spring Cankerworm, Gypsy Moth, Cabbage Looper, Imported Cabbage Worm, Tomato Hornworm



230 FM 87 • BONHAM, TEXAS 75418
EPA Reg. No. 70051-106-7401
EPA Est. No. 7401-TX-01
855-270-4776

16011-0922-REG

